

### TEST X (TOETS X) 3B

We use (Ons maak gebruik van)  $\{|0\rangle, |1\rangle\}$  as an orthonormal basis for a 2-dimensional Hilbert space in the following questions (vir 'n ortonormale basis vir 'n 2-dimensionele Hilbertruimte in die volgende vrae).

1) Calculate the following in terms of (Bereken die volgende in terme van)  $I, X, Y, Z$

a)  $XZ$

b)  $ZX$

c)  $U_{CNOT}(X \otimes I)U_{CNOT}$

d)  $U_{CNOT}(I \otimes X)U_{CNOT}$

e)  $U_{CNOT}(Z \otimes I)U_{CNOT}$

f)  $U_{CNOT}(I \otimes Z)U_{CNOT}$

g)  $U_{CNOT}(X \otimes X)U_{CNOT}$

h)  $U_{CNOT}(Z \otimes Z)U_{CNOT}$

i)  $U_{CNOT}U_{CNOT}$

where (waar)

$$I = |0\rangle\langle 0| + |1\rangle\langle 1|$$

$$X = |0\rangle\langle 1| + |1\rangle\langle 0|$$

$$Y = |0\rangle\langle 1| - |1\rangle\langle 0|$$

$$Z = |0\rangle\langle 0| - |1\rangle\langle 1|$$

2) Alice and Bob share  $n$  entangled pairs of the form (Alice en Bob deel  $n$  verstrikte pare)  $\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ . Thus we can write their shared state of  $2n$  qubits in the form

(Dus is dit moontlik om die gedeelde staat van die  $2n$  qubits soos volg te skryf)

$$|\psi\rangle = \frac{1}{\sqrt{2^n}} \sum_{j=0}^{2^n-1} |j\rangle \otimes |j\rangle$$

where the first  $n$  qubits belong to Alice and the second  $n$  qubits belong to Bob (waar die eerste  $n$  qubits aan Alice behoort, en die tweede  $n$  qubits aan Bob behoort). Furthermore Alice has  $2^n$  bits (Alice het  $2^n$  bisse)  $a_0, \dots, a_{2^n-1}$  and Bob has  $2^n$  bits (en Bob het  $2^n$  bisse)  $b_0, \dots, b_{2^n-1}$ . Calculate (Bereken)

$$|\phi\rangle := U_{PA} \otimes U_{PB} |\psi\rangle$$

$$\left( \bigotimes_n U_H \right) \otimes \left( \bigotimes_n U_H \right) |\phi\rangle.$$

where  $U_{PA}$  and  $U_{PB}$  acts on the computational basis as follows (waar  $U_{PA}$  en  $U_{PB}$  soos volg op die basis van berekening toegepas word)

$$U_{PA}|j\rangle = (-1)^{a_j}|j\rangle, \quad j = 0, 1, \dots, 2^n - 1$$

$$U_{PB}|j\rangle = (-1)^{b_j}|j\rangle, \quad j = 0, 1, \dots, 2^n - 1$$

For each of the cases (Vir die gevalle)

$$a) \quad a_0 = b_0, a_1 = b_1, \dots, a_{n-1} = b_{n-1}$$

$$b) \quad \sum_{k=0}^{2^n-1} |a_k - b_k| = 2^{n-1}$$

determine when measurement of the first  $n$  qubits in the computational basis yields the same result as measurement of the second  $n$  qubits in the computational basis (bepaal wanneer meting van die eerse  $n$  qubits in die basis van berekening presies dieselfde resultaat gee as meting van die tweede  $n$  qubits in die basis van berekening).